

BEHREND'S FUNCTION IS CONSTANT ON $\mathrm{Hilb}^n(\mathbb{C}^3)$.

ANDREW MORRISON

ABSTRACT. We prove that Behrend's function is constant on $\mathrm{Hilb}^n(\mathbb{C}^3)$. A calculation of motivic zeta functions shows the relevant Milnor fibers have zero Euler characteristic. As a corollary we see that $\mathrm{Hilb}^n(\mathbb{C}^3)$ is generically reduced. These results extend to moduli schemes of points and curves on resolutions of ADE singularities $\mathbb{C} \times Y_G$.

1. INTRODUCTION.

In [19] Thomas studies moduli spaces of sheaves on three dimensional Calabi–Yau manifolds and defines a holomorphic analogue of the Casson invariant associated to such moduli spaces. These integer valued invariants are deformation invariants and contain much interesting information about the geometry of the Calabi–Yau threefold in question [17] [15]. Here our philosophy is to consider the schemes $\mathrm{Hilb}^n(\mathbb{C}^3)$ as moduli spaces of stable sheaves on perhaps the simplest Calabi–Yau threefold, namely $\mathbb{C}^3$.

Some time after the introduction of these numbers it was realized by Behrend [1] that these Donaldson–Thomas invariants could be defined as Euler characteristics weighted by a certain constructible function.

Theorem 1.1. (*Behrend*) *For every scheme M over $\mathbb{C}$ there exists a canonical constructible function*

$$\nu_M : M \rightarrow \mathbb{Z}$$

such that if M is a projective moduli scheme with a symmetric obstruction theory then the associated Donaldson–Thomas type invariant is given by

$$\#^{vir}(M) = \sum_{n \in \mathbb{Z}} n \cdot \chi(\nu_M^{-1}(n)) \in \mathbb{Z}.$$

In this context we have a localization formula. Given a torus action with isolated fixed points this virtual count reduces to summing the values of ν_M at the fixed points. Behrend and Fantechi show that, for suitable torus actions, we have $\nu_M(p) = (-1)^{\dim T_p M}$ at each fixed point $p \in M$ [2]. In the case of $\mathrm{Hilb}^n(\mathbb{C}^3)$ there is a specific torus action whose fixed points correspond to three dimensional partitions. Moreover at these points $\nu_{\mathrm{Hilb}^n(\mathbb{C}^3)}(p) = (-1)^n$ [17] and we deduce that

$$\sum_{n \geq 0} \#^{vir}(\mathrm{Hilb}^n(\mathbb{C}^3)) t^n = \prod_{n \geq 1} (1 - (-t)^n)^{-n}$$

where the generating series on the right hand side is MacMahon's function enumerating three dimensional partitions, or the number of ways to stack n boxes in the corner of a room [14]. This gives the Donaldson–Thomas invariants for $\mathbb{C}^3$.

However the question still remains,

What are the values of Behrend's function elsewhere on $\text{Hilb}^n(\mathbb{C}^3)$?

Our main theorem provides a simple answer,

Theorem 1.2. *Behrend's function is constant with $\nu_{\text{Hilb}^n(\mathbb{C}^3)} = (-1)^n$.*

The proof of this theorem uses a relation to Milnor fibers. Indeed, if X is the scheme theoretic degeneracy locus of a regular function on a smooth variety $f : M \rightarrow \mathbb{C}$ then at $p \in X$

$$\nu_X(p) = (-1)^{\dim M} (1 - \chi(MF_{f,p})) \in \mathbb{Z}$$

where $MF_{f,p}$ is the Milnor fiber of f at p , i.e. the intersection of a nearby fibre of f with a small ball in M centred at p . Remarkably $\text{Hilb}^n(\mathbb{C}^3)$ has such a description as a critical locus. We can compute the relevant Milnor fibers, using arc spaces, and show that they have zero Euler characteristic. A complete proof of Theorem 1.2 is contained in Section 2.

It is more generally a fact that the value of Behrend's function at a point $p \in M$ is a local invariant of the singularity at p . Constancy of $\nu_{\text{Hilb}^n(\mathbb{C}^3)}$ puts restrictions on the singularities of $\text{Hilb}^n(\mathbb{C}^3)$. The following corollary to Theorem 1.2 is also contained in Section 2.

Corollary 1.3. *The components of $\text{Hilb}^n(\mathbb{C}^3)$ are generically reduced and have dimension equal to $n \bmod 2$.*

The remainder of the paper involves further calculations of motivic zeta functions. We prove a dimensional reduction result Theorem 5.3 in Section 5. In particular by applying this theorem to the case of $\text{Hilb}^n(\mathbb{C}^3)$ we see that the singularities of $\text{Hilb}^n(\mathbb{C}^3)$ are related to those of the commuting variety

$$\{(B, C) \in \text{End}(n)^{\times 2} : [B, C] = 0\}.$$

This leads to a second proof of Theorem 1.2 in Section 6. One advantage of the second proof is that it generalizes to other settings. In particular if Y_Q is a crepant resolution of an ADE singularity then $\mathbb{C} \times Y_Q$ is a Calabi-Yau threefold containing compact curves. Using Theorem 5.3 it is possible to show that Behrend's function is constant with absolute value equal to one on moduli schemes parameterizing points and curves in $\mathbb{C} \times Y_Q$. This theorem is stated here in the language of quiver moduli spaces (see Section 3).

Theorem 1.4. *Let Q, W be the quiver and potential associated to one of the ADE types. Then for θ -generic Behrend's function is constant on the moduli scheme of d -dimensional θ -stable framed $A = \mathbb{C}Q/I_W$ modules. Moreover we have*

$$\nu_{\{d\bar{f}_W=0\}} = (-1)^{\sum_{i \in I} f_i d_i}$$

where f_i are the components of the framing vector.

One new feature of this result is that it is valid in any chamber of the space of stability conditions. So for instance it is true in all of the PT/DT/ncDT chambers considered in generalized Donaldson–Thomas theory [13]. In particular, for the choice of stability condition corresponding to ideal sheaves we have the following.

Corollary 1.5. *For all numerical polynomials $p(n) = an + b$ the components of $\text{Hilb}^{p(n)}(\mathbb{C} \times Y_Q)$ are generically reduced and equi-dimensional mod 2.*

Theorem 1.4 and Corollary 1.5 are results that are peculiar to the geometry $\mathbb{C} \times Y_Q$. For moduli schemes parametrizing points and curves in general threefolds both constancy of the Behrend function and generic reducedness may fail.

Example 1.6. Non-Constancy. *This is achieved by the expository example of Pandharipande and Thomas [16] Section 4. They discuss degree 2 stable pairs with Euler number 4 on the resolved Conifold singularity. The moduli scheme of such objects has $\mathbb{P}^3$ as its underlying space but it contains a reduced scheme structure along a quadric hypersurface. Generically we have that Behrend's function equals -1 while along the quadric it takes value $+1$.*

Example 1.7. Non-Reducedness. *In [12] Section II, Mumford considers smooth degree 14 curves of genus 24 embedded in $\mathbb{P}^3$ and shows that over an open set in the Chow variety this family is incomplete. As a corollary of this we see that the Hilbert scheme of curves in $\mathbb{P}^3$ has a component that is generically non-reduced. This means that for Hilbert schemes of curves in threefolds not only is Behrend's function non-constant but it can have absolute value greater than one generically on an entire component.*

In Section 2 we prove Theorem 1.2. Section 3 and Section 4 contain background material on quiver moduli spaces and motivic zeta functions respectively. In Section 5 we prove the dimensional reduction statement Theorem 5.3. Section 6 applies Theorem 5.3 to relate the singularities of $\text{Hilb}^n(\mathbb{C}^3)$ to those of the commuting variety. Finally, we consider the ADE cases in Section 7.

2. PROOF OF MAIN THEOREM.

In this section we prove that Behrend's function is constant on $\text{Hilb}^n(\mathbb{C}^3)$. The three main properties of the function that we require are

- (1) (Locality) If $f : Y \rightarrow X$ is étale, then $f^*\nu_Y = \nu_X$,
- (2) (Multiplicativity) $\nu_{X \times Y}(p, q) = \nu_X(p) \cdot \nu_Y(q)$,
- (3) (Milnor Fiber) If X is the critical scheme of a regular function f on a smooth scheme M , i.e. $X = \{df = 0\}$, then

$$\nu_X(p) = (-1)^{\dim(M)}(1 - \chi(MF_{f,p}))$$

where $MF_{f,p}$ is the Milnor fiber of f at p .

The first of these implies that the value at a point is an invariant of the singularity. This allows to make calculations locally in a neighborhood of a given point. The second property means that if our scheme splits as a fiber product it is enough to find the values on each factor. While the third gives an explicit way to compute the value in the case of $\text{Hilb}^n(\mathbb{C}^3)$. All of these properties can be found in [1].

Theorem 1.2. *Behrend's function is constant with $\nu_{\text{Hilb}^n(\mathbb{C}^3)} = (-1)^n$.*

Proof. The first step is to setup the following description of $\text{Hilb}^n(\mathbb{C}^3)$ as coming from a quiver with potential,

$$\begin{array}{ccc} \text{GL}(V) & \longrightarrow & (\text{End}(V)^{\times 3} \times V)^{\text{stable}} \\ & & \downarrow \pi \\ & & (\text{End}(V)^{\times 3} \times V) //_{\chi} \text{GL}(V). \end{array} \quad \begin{array}{c} \xrightarrow{f} \mathbb{C} \\ \nearrow \tilde{f} \end{array}$$

Here V is an n -dimensional complex vector space. The group $\text{GL}(V)$ acts on the vector space V in the natural way and the quotient is given by Alistair King's construction of quiver moduli spaces [11]. In this case the stability condition is simply that the action of the three endomorphisms on the vector generates the entire vector space V . In particular the group action is free on the stable points and the map π is a principal $\text{GL}(V)$ bundle. The map f is defined by $f(A, B, C, v) = \text{Tr}(A[B, C])$. We have that

$$\text{Hilb}^n(\mathbb{C}^3) = \{d\tilde{f} = 0\}$$

is the scheme theoretic degeneracy locus of the regular function $\tilde{f}$ on the smooth space $(\text{End}(V)^{\times 3} \times V) //_{\chi} \text{GL}(V)$.

Basically, a point in this moduli space corresponds to a choice of three endomorphisms and a special vector satisfying the stability condition. The three endomorphisms are identified with the action of the co-ordinates x, y , and z on the ring of functions of our length n scheme while the special vector is identified with the unit in the ring, that the three endomorphisms lie in the degeneracy locus is precisely the further condition that they all commute. This construction is explained in detail in Example 3.2 of Section 3 where the general framework is developed for any quiver with potential.

The advantage of this description is that we can remove the action of the gauge group $\text{GL}(V)$ from all further considerations and work on the linear space of matrices in the pre-quotient. Locally, even Zariski locally, in a neighbourhood $U \subset \text{Hilb}^n(\mathbb{C}^3)$ the principal bundle $\pi : \{df = 0\} \rightarrow \text{Hilb}^n(\mathbb{C}^3)$ splits as $U \times \text{GL}(V)$. So at a point $(u, g) \in U \times \text{GL}(V) \subset \{df = 0\}$ we have

$$\begin{aligned} \nu_{\{df=0\}}(u, g) &= \nu_{\text{Hilb}^n(\mathbb{C}^3)}(u) \cdot \nu_{\text{GL}(V)}(g) \\ &= \nu_{\text{Hilb}^n(\mathbb{C}^3)}(u) \cdot (-1)^{n^2}. \end{aligned}$$

Here the first equality follows from the multiplicativity property and the second is a consequence of the fact that on a smooth scheme Y , $\nu_Y(y) = (-1)^{\dim(Y)}$ for all $y \in Y$ [1]. As mentioned it now suffices to work on the prequotient and prove the equivalent statement that $\nu_{\{df=0\}}$ is constant with value equal to 1. We are reduced to studying the critical locus of the function $f : \text{End}(V)^{\times 3} \times V \rightarrow \mathbb{C}$ given by $f((A, B, C, v)) = \text{Tr}(A[B, C])$ and we can use the relation to Milnor fibres.

- (3) (Milnor Fiber) If X is the critical scheme of a regular function f on a smooth scheme M , i.e. $X = \{df = 0\}$, then

$$\nu_X(p) = (-1)^{\dim(M)}(1 - \chi(MF_{f,p}))$$

where $MF_{f,p}$ is the Milnor fiber of f at p .

Since in our specific case the dimension of our ambient linear space has even dimension equal to $3n^2 + n$ coming from the three endomorphisms and a vector we see that the value of Behrend's function is constant with value equal to 1 if and only if

$$\chi(MF_{f,p}) = 0$$

for all p in the critical set $\{df = 0\}$. We will attack this problem directly using arc spaces. Arc spaces are defined functorially with the underling set of $\mathbb{C}$ -rational points determined by maps of a curvilinear fat point into our variety

$$\mathcal{L}_n(X)(\mathbb{C}) := \text{Hom}(\text{Spec}(\mathbb{C}[t]/(t^{n+1})), X)$$

moreover if we insist that the topological support of the arc be located at a point $p \in X$, we can study the space of arcs based at p , these spaces can record interesting information about the singularities of X at p

$$\mathcal{L}_n(p, X)(\mathbb{C}) := \{\gamma(t) \in \mathcal{L}_n(X) : \text{supp}(\gamma(t)) = p\}.$$

Now given a regular function $f : X \rightarrow \mathbb{A}_{\mathbb{C}}^1$ by the functoriality of the construction we get a map $f : \mathcal{L}_n(X) \rightarrow \mathcal{L}_n(\mathbb{A}_{\mathbb{C}}^1)$ sending $\gamma(t)$ to $f \circ \gamma(t)$. In coordinates this arc in $\mathbb{A}_{\mathbb{C}}^1$ can be expressed as

$$f \circ \gamma(t) = a_0 + a_1 t + \cdots a_n t^n \mod t^{n+1}.$$

We define

$$\mathcal{X}_n(p) = \{\gamma \in \mathcal{L}_n(p, X) : f \circ \gamma(t) = t^n\}$$

these are arcs in X based at p that vanish up to order $n - 1$ into the central fiber. These spaces provide a measure of the singularities of f . Indeed the zeta function

$$Z_{f,p}(T) := \sum_{n \geq 1} \chi(\mathcal{X}_n(p)) T^n \in \mathbb{Z}[[T]]$$

is known to be a rational function with removable singularity at ∞ such that $\lim_{T \rightarrow \infty} Z_{f,p}(T) = \chi(MF_{f,p})$ [4]. In our case we are considering the function $f : \text{End}(V)^{\times 3} \times V \rightarrow \mathbb{C}$ given by $f(A, B, C, v) = \text{Tr}(A[B, C])$. From now on we fix a point $p = (A_0, B_0, C_0, v_0)$ in the critical locus of f at which we are interested in computing the Milnor fibre $MF_{f,p}$. This corresponds to a choice of a vector and three commuting matrices. The arc spaces $\mathcal{X}_n(p)$ are given by

$$\left\{ (A_i, B_i, C_i, v_i)_{i=1 \dots n} : \text{Tr} \left(\sum_{i=0}^n A_i t^i \left[\sum_{i=0}^n B_i t^i, \sum_{i=0}^n C_i t^i \right] \right) = t^n \mod t^{n+1} \right\}.$$

We are going to show that all of these spaces have Euler charateric equal to zero. This will imply

$$\chi(\mathcal{X}_n(p)) = 0 \Rightarrow Z_{f,p}(T) = 0 \Rightarrow \chi(MF_{f,p}) = 0$$

giving the result. To this end we consider the group of matrices

$$G_{A,B,C} := \{P \in \text{GL}(V) : [P, A] = [P, B] = [P, C] = 0\}$$

this group acts on the set $\mathcal{X}_n(p)$ via $g(A_i, B_i, C_i, v_i) = (gA_i g^{-1}, gB_i g^{-1}, gC_i g^{-1}, gv_i)$. Moreover the group has Euler characteristic equal to zero and so

$$\chi(\mathcal{X}_n(p)) = \chi(\mathcal{X}_n(p))^{G_{A,B,C}}.$$

We turn our attention to the fixed points of this action. These are given precisely by

$$gA_i g^{-1} = A_i, gB_i g^{-1} = B_i, gC_i g^{-1} = C_i, \text{ and } v_i = 0.$$

So that the matrices A_i, B_i, C_i commute with every element in the group $G_{A,B,C}$. This implies that every matrix present commutes with A_0, B_0, C_0 . In fact (although it is unnecessary) without loss of generality we could have chosen the three matrices A_0, B_0, C_0 to have been invertible from the start and hence present in the group $G_{A,B,C}$ by fiat. This commutativity implies that the trace function has a simplified form when restricted to the fixed point set $\mathcal{X}_n(p)^{G_{A,B,C}}$ given by

$$\mathrm{Tr} \left(\sum_{i=1}^n A_i t^i \left[\sum_{i=1}^n B_i t^i, \sum_{i=1}^n C_i t^i \right] \right) = t^n \pmod{t^{n+1}}.$$

Now all sums begin at $i = 1$ and the polynomial is homogeneous in the variables. So we have a $\mathbb{C}^*$ action on the fixed point set given by $\lambda(A_i, B_i, C_i) = (\lambda^2 A_i, \lambda^{-1} B_i, \lambda^{-1} C_i)$. This action is free so that

$$\chi(\mathcal{X}_n(p)^{G_{A,B,C}}) = \chi((\mathcal{X}_n(p)^{G_{A,B,C}})^{\mathbb{C}^*}) = \chi(\emptyset)$$

which equals zero. $\square$

Corollary 1.3. *The components of $\mathrm{Hilb}^n(\mathbb{C}^3)$ are generically reduced and have dimension equal to $n \bmod 2$.*

Proof. Suppose X is an irreducible scheme then at a general point $p \in X$ the value of Behrend's function is given by

$$\nu_X(p) = (-1)^{\dim(X)} \cdot m$$

where $m \in \mathbb{N}$ is the generic multiplicity X . Indeed, on X the value of ν_X can be defined cycle theoretically via a signed version of MacPherson's local Euler obstruction on the normal cone ([7] Chapter 4.3). Now the corollary follows from our theorem by applying the above fact to each of the components of $\mathrm{Hilb}^n(\mathbb{C}^3)$. $\square$

Remark 2.1. *Given any smooth threefold X . Then étale locally $\mathrm{Hilb}^n(X)$ is isomorphic to*

$$\mathrm{Hilb}^{n_1}(\mathbb{C}^3) \times \cdots \times \mathrm{Hilb}^{n_k}(\mathbb{C}^3)$$

for some $n_1 + \cdots + n_k = n$ [8]. From this the above theorem and corollary actually extend to any smooth threefold using the locality and multiplicativity properties (1) and (2) above.

3. QUIVER MODULI SPACES.

Here we develop a general framework of quiver moduli spaces. This includes the above description of $\mathrm{Hilb}^n(\mathbb{C}^3)$ (see Example 3.2). The reader only interested in these schemes may wish to skip this material. The construction will mainly be used for the ADE resolutions in Section 7.

Let $Q = (i \in I, \alpha : i \rightarrow j)$ be a finite dimensional quiver. A representation of this quiver is given by $M = (V_i, M_\alpha : V_i \rightarrow V_j)$ a collection of vector spaces V_i one based at each vertex and a linear map $M_\alpha : V_i \rightarrow V_j$ associated to each edge. These representations naturally form an abelian category $\mathrm{rep}(Q)$. Every representation has a dimension vector $\dim(M) = (\dim(V_i))_{i \in I} \in \mathbb{Z}^I$. In fact these categories are hereditary and if we define a bilinear pairing on $\mathbb{Z}^I$ via,

$$\langle d, e \rangle_Q = \sum_{i \in I} d_i e_i - \sum_{\alpha : i \rightarrow j} d_i e_j$$

then we have

$$\chi(M, N) = \text{Ext}^0(M, N) - \text{Ext}^1(M, N) = \langle \dim(M), \dim(N) \rangle_Q.$$

The connection of these representations to algebra is to observe that we can define a path algebra $\mathbb{C}Q$, with a vector space basis given by paths in the quiver, and multiplication given by concatenation of such subsequent paths. This gives an isomorphism between the category of finite dimensional representation of the quiver and finite dimensional modules for the path algebra, $\text{rep}(Q) = \text{mod}(\mathbb{C}Q)$. Considering these objects up to isomorphism we have the moduli stack of Q -representations of dimension $d \in \mathbb{Z}$

$$\mathfrak{M}_d(Q) := [H_d/G_d]$$

with $H_d := \prod_{\alpha:i \rightarrow j} \text{Hom}(\mathbb{C}^{d_i}, \mathbb{C}^{d_j})$ and $G_d := \prod_{i \in I} \text{GL}(\mathbb{C}^{d_i})$ and where the group G_d acts on H_d in the natural way. In general the stabilizer groups in this quotient will be affine algebraic groups of positive dimension. We would like to have a smooth quotient in the category of schemes Alistair King's idea [11] was to use Mumford's projective GIT. Given $\theta \in (\mathbb{Q}^*)^I$ we say that a d dimensional representation M is θ -(semi-)stable if

$$\mu(N) := \frac{\theta(e)}{\sum_{i \in I} e_i} (\leq) < \mu(M) := \frac{\theta(d)}{\sum_{i \in I} d_i}$$

for all $0 \neq N \subsetneq M$ with $\dim(N) = e$. Furthermore to ensure the moduli spaces are smooth we introduce a framing vector $f \in \mathbb{Z}_{>0}^I$ and study framed representations. These consist of a representation of the original quiver and a collection of f_i vectors in each the vector spaces V_i . In [6] a modified stability condition is introduced, we say that a framed representation (M, v) is θ -stable if

$$M \text{ is } \theta \text{ semistable and } \mu(U) < \mu(M)$$

for any proper subrepresentation U containing the framing vectors v_i . If we define

$$M_d^{f, \theta}(Q) = (H_d \times V_f)^{\theta \text{ stable}} / G_d$$

to be the geometric quotient of the set of θ -stable framed representations. Then it is proven [6].

Theorem 3.1. *For fixed d and θ -generic, we have that $M_d^{f, \theta}(Q)$ is a smooth fine moduli space parameterizing stable framed quiver representations of dimension d . In particular, the quotient map*

$$\pi : (H_d \times V_f)^{\theta \text{ stable}} \rightarrow M_d^{f, \theta}(Q)$$

is a principal G_d bundle.

Here θ -generic means that theta is chosen so that stability coincides with semi-stability for all (non-framed) representations of dimension d . Indeed this is an open condition on the set of stability parameters.

Now that we have produced these smooth moduli spaces we want to consider a potential function on them, the critical locus of which we will be interested in.

A potential is an element of the path algebra $W \in \mathbb{C}Q$ such that W is a formal linear combination of paths that are cycles. We can use this to define a polynomial f_W in the following way

$$f_W : H_d \times V_f \rightarrow \mathbb{C} \text{ via } f_W(M, v) = \text{Tr}(M_\alpha),$$

by taking the sum of the traces around each cycle. The function is clearly equivariant for the gauge group G_d and induces a regular function

$$\overline{f}_W : M_d^{f,\theta}(Q) \rightarrow \mathbb{C}$$

the critical locus of such a function carries a symmetric obstruction theory and led to Szendroi's [18] definition of non-commutative Donaldson-Thomas invariants. The name coming from the fact that the critical locus of this function parameterises modules for the (possibly non-commutative) three dimensional Calabi-Yau algebra

$$A = \mathbb{C}Q/I_W$$

where I_W is the two sided ideal generated by the cyclic derivatives of the potential [8].

Example 3.2. *The Hilbert scheme of points on $\mathbb{C}^3$. We first consider the 3 loop quiver $Q_3 = (\star, \{x, y, z\})$ with one vertex and three arrows. The path algebra for this quiver with multiplication given by concatenation of paths is clearly seen to be equal to $\mathbb{C}\langle x, y, z \rangle$. We will take $\theta = 1 \in \mathbb{Q}^*$ and a single framing vector $f = 1 \in \mathbb{Z}$. The smooth moduli space*

$$M_n^{1,1}(Q_3)$$

then parasitizes n -dimensional cyclic modules for $\mathbb{C}\langle x, y, z \rangle$ or equivalently left ideals of length n . Next we take the potential $W = x(yz - zy)$, then the vanishing of df_W implies commutation relations for the maps in the representation, as an exercise one can check

$$A = \mathbb{C}Q/I_W = \mathbb{C}[x, y, z].$$

The scheme theoretic degeneracy locus of this potential is

$$\text{Hilb}^n(\mathbb{C}^3) = \{df_W = 0\} \subset M_n^{1,1}(Q_3)$$

the Hilbert scheme of n points in $\mathbb{C}^3$. This description was used prominently in [3].

Remark 3.3. *In the above example there are interesting non-commutative deformations most notably the Sklyanin algebras [20] given by modifying this potential to*

$$W = a \cdot xyz + b \cdot xzy + \frac{c}{3} (x^3 + y^3 + z^3)$$

where $[a : b : c] \in \mathbb{P}^2$. We can extend the results of this paper to the cases when $c = 0$ but the other cases remain interesting.

4. MOTIVIC ZETA FUNCTIONS.

Motivic zeta functions were introduced by Denef and Loeser [4] as a geometric analogue of the p -adic zeta functions studied by Igusa [10]. We will recall here their definition and some of their properties relevant for our purposes.

Let S be a scheme over $\mathbb{C}$. We define the Grothendieck group of S -varieties $K_0(\text{Var}_S)$ to be the free abelian group on isomorphism classes of S -varieties modulo the scissor relations,

$$[X \rightarrow S] = [Z \rightarrow S] + [X \setminus Z \rightarrow S]$$

for any $Z \subset X$ a subvariety. This group can be made into a ring using the fibre product as multiplication

$$[X \rightarrow S] \times [Y \rightarrow S] = [X \times_S Y \rightarrow S].$$

We will need to make one slight extension of this ring to varieties with a group action. Let G be a group with a good action on S , we say that an action is good if it preserves a collection of affine charts. Then we can define $K^G(\text{Var}_S)$ to be the collection of G -equivariant S -varieties. That is isomorphism classes of G -equivariant maps $[X \rightarrow S]$ where X has also a good G action. Again we impose the scissor relations and the obvious fiber product. However in this ring it is very convenient to quotient out by the further relations,

$$[V \times_{\mathbb{C}} S \rightarrow S] = [\mathbb{A}_{\mathbb{C}}^{\dim(V)} \times_{\mathbb{C}} S \rightarrow S]$$

where V is any G -representation and the class on the right has the trivial action. So we make these extra assumptions and as a piece of notation write $\mathbb{L} = [\mathbb{A}_{\mathbb{C}}^1 \times_{\mathbb{C}} S \rightarrow S]$ for the class of the affine line. These classes we will often refer to as motivic classes, they can be thought of as universal Euler characteristics. Indeed there is a map

$$\chi : K^G(\text{Var}_S) \rightarrow \mathbb{Z} \text{ given by } [X] \mapsto \chi(X)$$

from the Grothendieck group of S -varieties to the integers given by taking the numerical Euler characteristic. Indeed this is a homomorphism of rings if $S = \text{Spec}(\mathbb{C})$ is a point. Similar maps exist for the E -polynomial, Poincare polynomial, etc [4].

In order to define the motivic zeta function we also need to make a brief discussion of arc spaces (a.k.a. jet schemes). We give a modular definition, by defining the n th arc space of the complex scheme X to be the separated scheme $\mathcal{L}_n(X)$ representing the functor,

$$F_n : (\mathbb{C} - \text{algs}) \rightarrow (\text{Sets}) \text{ given by } R \mapsto \text{Hom}(\text{Spec}(R[t]/(t^{n+1}), X)).$$

In fact it is easy to prove that the above functor is representable using local coordinates for X . Notice that truncation of the curvilinear schemes gives natural maps $\pi_n^{n+1} : \mathcal{L}_{n+1}(X) \rightarrow \mathcal{L}_n(X)$. Moreover by the functionality of the construction given any map $f : X \rightarrow Y$ we get an induced map $\mathcal{L}_n(f) : \mathcal{L}_n(X) \rightarrow \mathcal{L}_n(Y)$, given by sending $\gamma \in \mathcal{L}_n(X) \rightarrow f \circ \gamma \in \mathcal{L}_n(Y)$. In this way arc spaces can be seen as a generalization of the tangent bundle construction.

Now suppose that we have a regular function $f : X \rightarrow \mathbb{C}$ on a smooth space X . Let $X_0 = f^{-1}(0)$, we can define the following arc spaces in the hope of measuring the singularities of X_0

$$\mathfrak{X}_n := \{\gamma \in \mathcal{L}_n(X) : f \circ \gamma(t) = a_n t^n \mod t^{n+1} \text{ with } a_n \in \mathbb{C}^*\}$$

these are n -arcs that vanish to order $n - 1$ in X_0 . This space has a natural $\mathbb{C}^*$ action given by $z \cdot \gamma(t) = \gamma(z \cdot t)$ for $z \in \mathbb{C}^*$. Moreover they are relative over the central fibre $X_0 = f^{-1}(0)$. We can define the naive motivic zeta function

$$Z_f^{\text{naive}}(T) := \sum_{n \geq 1} [\mathfrak{X}_n \rightarrow X_0] (\mathbb{L}^{-\dim(X)} T)^n \in K_0(\text{Var}/X_0)[\mathbb{L}^{-1}][[T]].$$

Now consider the subspaces

$$\mathfrak{X}_{n,1} := \{\gamma \in \mathcal{L}_n(X) : f \circ \gamma(t) = t^n \mod t^{n+1}\} \subset \mathfrak{X}_n.$$

The above $\mathbb{C}^*$ action gives a restricted μ_n action on these spaces. If we define $\hat{\mu} = \lim \mu_n$ to be the protective limit of these groups then we can define the motivic

zeta function

$$Z_f(T) := \sum_{n \geq 1} [\mathfrak{X}_{n,1} \rightarrow X_0, \mu_n](\mathbb{L}^{-\dim(X)} T)^n \in K^{\hat{\mu}}(\text{Var}/X_0)[\mathbb{L}^{-1}][[T]].$$

Finally as a conglomeration of these two functions we define

$$Z_f^{\phi}(T) := (-1)^{\dim(X)} \left(\frac{T[X_0] - Z_f^{\text{naive}}(T)}{1 - T} - Z_f(T) \right) \in K^{\hat{\mu}}(\text{Var}/X_0)[\mathbb{L}^{-1}][[T]].$$

A theorem of Denef and Loeser says that all of these zeta functions are rational functions and moreover the limit as $T \rightarrow \infty$ exists.

Theorem 4.1 (Theorem 3.3.1 [4]). *The functions defined above are rational functions in T moreover they can be written as a finite sum of functions of the form*

$$a \cdot \prod_{i=1}^k \frac{\mathbb{L}^{-\nu_i} T^{N_i}}{1 - \mathbb{L}^{-\nu_i} T^{N_i}}$$

where $a \in K^{\hat{\mu}}(\text{Var}/X_0)$ and $\nu_i, N_i \in \mathbb{N}$.

In fact their theorem says much more giving an explicit formula for a rational function in terms of an embedded resolution, but we will not take this approach in our computations. Now we can define the motivic vanishing cycle of $f : X \rightarrow \mathbb{C}$

$$[\phi_f] := \lim_{T \rightarrow \infty} Z_f^{\phi}(T) \in K^{\hat{\mu}}(\text{Var}/X_0).$$

Remark 4.2. *All of the definitions given above were relative over the central fibre. Choosing a point $p \in X_0$ we can make all of these constructions relative to a point by pulling back all the motivic classes along the map $i_p : p \rightarrow X_0$ we get*

$$\begin{aligned} i_p^* &: K^{\hat{\mu}}(\text{Var}/X_0) \rightarrow K^{\hat{\mu}}(\text{Var}/p) \\ &: [\mathcal{X} \rightarrow X_0] \mapsto [p \times_{X_0} \mathcal{X} \rightarrow p] \end{aligned}$$

for $p \in X_0$. For example we can define a motivic zeta function $Z_{f,p}(T)$ relative to a point $p \in X_0$ in terms of arcs spaces

$$\mathcal{X}_n(p) := \{\gamma \in \mathcal{X}_n : \text{supp}(\gamma) = p\}$$

i.e. arcs based at the point p . Then we have,

$$[\phi_{f,p}] := i_p^*[\phi_f] = \lim_{T \rightarrow \infty} Z_{f,p}^{\phi}(T).$$

It is again a theorem (see Theorem 3.5.5 [4]) that this motivic vanishing cycle has all the usual properties of the sheaf of vanishing cycles. In particular

$$\chi([\phi_{f,p}]) = (-1)^{\dim(X)} (1 - \chi(MF_{f,p})) \in \mathbb{Z}$$

where $MF_{f,p}$ is the analytic Milnor fibre. This completes our general discussion of motivic zeta functions.

We now turn to a special case studied by Davison and Meinhardt [5]. Henceforth we consider a function $f : \mathbb{A}^d \rightarrow \mathbb{C}$. Moreover we assume there exists a $\mathbb{C}^*$ action on $\mathbb{A}^d$ given by $t(x_1, \dots, x_d) = (tx_1, \dots, tx_r, x_{r+1}, \dots, x_d)$ such that for this action we have $f(t \cdot x) = t^p f(x)$ for some $p \in \mathbb{N}$. Let $\mathcal{L}_n(\mathbb{A}^d)^0 \subset \mathcal{L}_n(\mathbb{A}^d)$ be the collection of arcs with topological support in $X_0 = f^{-1}(0)$. Then we consider the composite map

$$f_n : \mathcal{L}_n(\mathbb{A}^d)^0 \xrightarrow{\mathcal{L}_n(f)} \mathcal{L}_n(\mathbb{A}^1) \longrightarrow \mathbb{A}^1$$

given by

$$f_n : \gamma(t) \mapsto f \circ \gamma(t) = a_1 t + \dots + a_n t^n \mapsto a_n.$$

The fibres of f_n over zero and one both have μ_n actions analogous to those above and it is shown ([5] Section 5)

$$Z_f^\phi(T) = (-1)^d \sum_{n \geq 1} ([f_n^{-1}(0) \rightarrow X_0, \mu_n] - [f_n^{-1}(1) \rightarrow X_0, \mu_n]) (\mathbb{L}^{-d} T)^n$$

as a power series in the ring $K^{\hat{\mu}}(\text{Var}/X_0)[\mathbb{L}^{-1}][[T]]$. In fact by using the further relations on the $\hat{\mu}$ equivariant motives above it is shown that one can take the action on $[f_n^{-1}(0)]$ to be trivial ([5] Remark 4.2).

Remark 4.3. *On the level of Euler numbers it is possible to show that this formula is true regardless of the equivariance assumptions on f . Indeed at a point $p \in X_0$ we always have*

$$\chi\left(Z_{f,p}^\phi(T)\right) = (-1)^d \sum_{n \geq 1} (\chi(f_n^{-1}(0) \times_{X_0} p) - \chi(f_n^{-1}(1) \times_{X_0} p)) T^n \in \mathbb{Z}[[T]]$$

for any function $f : \mathbb{A}^d \rightarrow \mathbb{C}$.

5. DIMENSIONAL REDUCTION.

In this section we discuss a method that can be used to compute $Z_{f,p}^\phi(T)$ under some very restrictive assumptions on the function f .

Definition 5.1. *Let $(\mathbb{A}^l, \langle, \rangle)$ be a complex vector space endowed with a non-degenerate bilinear form and suppose there exists a homogenous function $r : \mathbb{A}^r \rightarrow \mathbb{A}^l$ such that,*

$$\begin{aligned} f & : \mathbb{A}^l \times \mathbb{A}^r \rightarrow \mathbb{C} \\ & : (x, y) \mapsto \langle x, r(y) \rangle, \end{aligned}$$

then we say that f has a linear factor.

Definition 5.2. *Associated to a function with linear factor as above. We define the reduced variety to be*

$$R = \{y \in \mathbb{A}^r : r(y) = 0\}.$$

The dimensional reduction theorem we shall prove relates the motivic zeta function of f to the arc spaces of this reduced variety. Throughout this discussion we will be interested in relative calculations and as such we fix once and for all a point $p = (x_0, y_0) \in \{df = 0\} \subset X_0$. Notice that in particular this implies that $r(y_0) = 0$.

We consider the following collection of arcs based at y_0

$${}^n R := \left\{ (y_i)_{i=1, \dots, n} \in \mathbb{A}^{r \cdot n} : r \left(\sum_{i=0}^{n-1} y_i t^i \right) = 0 \pmod{t^n} \right\}.$$

We can regard this as a sub-variety of the arc space $\mathcal{L}_n(\mathbb{A}^r, y_0)$ as such it is isomorphic to $\mathcal{L}_{n-1}(y_0, R) \times \mathbb{A}^r$, given by the arc space of the reduced variety at the point y_0 . Then we can consider the space $\mathbb{A}^{l \cdot n} \times ({}^n R)$ as a subspace of

$\mathcal{L}_n(\mathbb{A}^l, (x_0)) \times \mathcal{L}_n(\mathbb{A}^r, (y_0)) = \mathcal{L}_n(\mathbb{A}^{l+r}, (x_0, y_0))$. On restricting to this sub-variety the function f_n defined above takes a much simpler form

$$f_n|_{\mathbb{A}^{l \cdot n} \times ({}^n R)} : (x_i, y_i)_{i=1, \dots, n} \mapsto \langle x_0, r \left(\sum_{i=1}^n y_i t^i \right) \rangle \mod t^{n+1}$$

in particular the function has no dependence on the x_i . Motivated by this we define

$$\begin{aligned} \Lambda_n^0(x_0) &:= \left\{ (y_i) \in ({}^n R) : \langle x_0, r \left(\sum_{i=0}^n y_i t^i \right) \rangle = 0 \mod t^{n+1} \right\} \\ \Lambda_n^1(x_0) &:= \left\{ (y_i) \in ({}^n R) : \langle x_0, r \left(\sum_{i=0}^n y_i t^i \right) \rangle = t^n \mod t^{n+1} \right\} \end{aligned}$$

where both the spaces above have the natural μ_n action given by $\zeta \cdot (y_i) = (\zeta^i y_i)$ for a primitive n th root ζ .

Theorem 5.3. *We have*

$$Z_{f,p}^\phi(T) = (-1)^{l+r} \sum_{n \geq 1} ([\Lambda_n^0(x_0), \mu_1] - [\Lambda_n^1(x_0)]) (\mathbb{L}^{-r} T)^n \in K^{\hat{\mu}}(Var/p)[\mathbb{L}^{-1}][[T]].$$

Proof. First we remark that any function f with a linear factor satisfies the equivariance condition at the end of our discussion on motivic zeta functions. In fact we have a $\mathbb{C}^*$ action on $\mathbb{A}^{l+r}$ given by $t(x, y) = (tx, y)$ such that $f(t(x, y)) = tf(x, y)$. Therefore as above we can write

$$Z_{f,p}^\phi(T) = (-1)^{l+r} \sum_{n \geq 1} ([f_n^{-1}(0) \times_{X_0} p, \mu_n] - [f_n^{-1}(1) \times_{X_0} p, \mu_n]) (\mathbb{L}^{-l-r} T)^n$$

where to make the function relative to the point $p = (x_0, y_0)$ we simply pulled back along the inclusion map $i_p : p \rightarrow X_0$ (see Remark 4.2). The rest on the proof just involves cancelation of certain terms at each coefficient T^n .

The function f_n is defined on the space of arcs based at p via

$$\begin{aligned} f_n &: \mathcal{L}_n(\mathbb{A}^d) \rightarrow \mathbb{C} \\ &: (x_i, y_i) \mapsto \left\langle \sum_{i=0}^n x_i t^i, r \left(\sum_{i=0}^n y_i t^i \right) \right\rangle \mod t^{n+1}. \end{aligned}$$

We define

$${}^n R^m := \left\{ (y_i) \in \mathbb{A}^{r \cdot n} : r \left(\sum_{i=0}^n y_i t^i \right) = v_m t^m + v_{m+1} t^{m+1} \dots \mod t^{n+1}, v_m \neq 0 \right\}.$$

These spaces provide a μ_n equivariant stratification of the space of arcs

$$\mathcal{L}_n((x_0, y_0), \mathbb{A}^l \times \mathbb{A}^r) = \mathbb{A}^{ln} \times ({}^n R^1) \sqcup \dots \sqcup \mathbb{A}^{ln} \times ({}^n R^n)$$

notice that ${}^n R^n$ is just ${}^n R$ defined above. Now when restricted to each of these components f_n takes a simplified form

$$\begin{aligned} f_n &: \mathbb{A}^{nl} \times ({}^n R^m) \rightarrow \mathbb{C} \\ &: (x_i, y_i) \mapsto \langle x_{n-m}, v_m \rangle + \sum_{i \geq m} \langle x_{n-i}, v_i \rangle. \end{aligned}$$

Let us assume that $m < n$. If we insist that $f_n((x_i, y_i)) = 1$ then the condition that $v_m \neq 0$ imposes a single linear condition on the vector x_{n-i} while all other x_i coordinates are unrestricted, we get that

$$[f_n|_{\mathbb{A}^{nl} \times ({}^n R^m)}(1)] = \mathbb{L}^{n^{l-1}}[{}^n R^m, \mu_n].$$

Similarly, an identical formula holds for the inverse image of zero over $\mathbb{A}^{nl} \times ({}^n R^m)$. This gives the desired cancelation

$$[f_n^{-1}(0)] - [f_n^{-1}(1)] = [f_n|_{\mathbb{A}^{nl} \times ({}^n R)}(0)] - [f_n|_{\mathbb{A}^{nl} \times ({}^n R)}(1)].$$

We saw above in our discussion prior to the theorem that the function f_n restricted to $\mathbb{A}^{nl} \times ({}^n R)$ bears no dependence on the coordinates x_i . From this it follows immediately that

$$\begin{aligned} [f_n|_{\mathbb{A}^{nl} \times ({}^n R)}(0)] &= \mathbb{L}^{nl}[\Lambda_n^0(x_0)] \\ [f_n|_{\mathbb{A}^{nl} \times ({}^n R)}(1)] &= \mathbb{L}^{nl}[\Lambda_n^1(x_0)] \end{aligned}$$

and both these motives inherit the μ_n action described above. Moreover we may take the action on $\Lambda_n^0(x_0)$ to be trivial as mentioned in [5] Remark 4.2. $\square$

Corollary 5.4. *Let $f : \mathbb{A}^{l+r} \rightarrow \mathbb{C}$ be a function with a linear factor as above and let $p \in \{df = 0\} \subset \mathbb{A}^{l+r}$ then*

$$\nu_{\{df=0\}}(p) = \lim_{T \rightarrow \infty} \chi(Z_{f,p}(T))$$

where

$$\chi(Z_{f,p}(T)) = (-1)^{l+r} \sum_{n \geq 1} (\chi(\Lambda_n^0(x_0)) - \chi(\Lambda_n^1(x_0))) T^n.$$

Proof. This corollary is an immediate consequence of [4] Theorem 3.5.5 mentioned in Section 3 and the Milnor fibre description of ν [1] given by

$$\nu_{\{df=0\}}(p) = (-1)^{l+r} (1 - \chi(MF_{f,p})).$$

$\square$

Remark 5.5. *The above corollary remains true if we drop the condition that the function $r : \mathbb{A}^r \rightarrow \mathbb{A}^l$ is homogeneous. We will not need this here but the proof of Theorem 5.3 will carry over in this setting by neglecting the μ_n actions and comparing with Remark 4.3.*

6. SINGULARITIES AND THE COMMUTING VARIETY.

In this section we are going to use the dimensional reduction theorem above to give a second argument that the Milnor fibres of

$$\begin{aligned} f &: \text{End}(V)^{\times 3} \rightarrow \mathbb{C} \\ &: (A, B, C) \mapsto \text{Tr}(A[B, C]) \end{aligned}$$

are equal to zero. Therefore implying the constancy of Behrend's function on $\text{Hilb}^n(\mathbb{C}^3)$.

It is clear that the map f has a linear factor, where we consider the trace form as a non-degenerate pairing on $\text{End}(V)$. The corresponding reduced variety equals

$$R = \{(B, C) \in \text{End}(V)^{\times 2} : [B, C] = 0\}$$

and is known as the commuting variety. At a point $p = (A_0, B_0, C_0) \in \{df = 0\}$ corresponding to three commuting matrices we have

$$\begin{aligned}\Lambda_n^0(A_0) &= \{(B_i, C_i)_{i=1, \dots, n-1} \in \mathcal{L}_{n-1}(R, (B_0, C_0)) \\ &\quad : \text{Tr} \left(A_0, \left[\sum_{i=1}^{n-1} B_i t^i, \sum_{i=1}^{n-1} C_i t^i \right] \right) = 0 \pmod{t^{n+1}} \} \\ &\quad \times \text{End}(V)^{\times 2}.\end{aligned}$$

$$\begin{aligned}\Lambda_n^1(A_0) &= \{(B_i, C_i, v_i)_{i=1, \dots, n-1} \in \mathcal{L}_{n-1}(R, (B_0, C_0)) \\ &\quad : \text{Tr} \left(A_0, \left[\sum_{i=1}^{n-1} B_i t^i, \sum_{i=1}^{n-1} C_i t^i \right] \right) = t^n \pmod{t^{n+1}} \} \\ &\quad \times \text{End}(V)^{\times 2}.\end{aligned}$$

Here we can consider the sums beginning at one by the commutativity of the three matrices A_0, B_0, C_0 . Now by the results of the last section

$$Z_{f,p}^\phi(T) = (-1)^{\dim(V)^3} \sum_{n \geq 1} ([\Lambda_n^0(A_0), \mu_1] - [\Lambda_n^1(A_0), \mu_n]) (\mathbb{L}^{-2 \dim(V)^2} T)^n$$

and in particular

$$\chi(Z_{f,p}^\phi(T)) = (-1)^{\dim(V)^3} \sum_{n \geq 1} (\chi(\Lambda_n^0(A_0)) - \chi(\Lambda_n^1(A_0))) T^n.$$

Then we can consider the action of $\mathbb{C}^*$ given by $\lambda \cdot (B_i, C_i) = (\lambda B_i, \lambda^{-1} C_i)$ preserving both $\Lambda_n^1(A_0)$ and $\Lambda_n^0(A_0)$. The action on the first space is free while the action on the second has a unique fixed point $(0, 0)$, this gives

$$\chi(\Lambda_n^1(A_0)) = 0 \text{ and } \chi(\Lambda_n^0(A_0)) = 1.$$

Taking $\lim_{T \rightarrow \infty} \chi(Z_{f,p}(T))$ implies all Milnor fibers have zero Euler characteristic.

7. ADE RESOLUTIONS.

Lastly we look at some other threefolds for which the value of Behrend's function remains constant. Consider the Kleinian surface singularities given by

$$\begin{aligned}A_n &: w^2 + x^2 + y^{n+1} = 0 \\ D_n &: w^2 + y(x^2 + y^{n-2}) = 0 \quad (n \geq 4) \\ E_6 &: w^2 + x^3 + y^4 = 0 \\ E_7 &: w^2 + x(x^2 + y^3) = 0 \\ E_8 &: w^2 + x^3 + y^5 = 0.\end{aligned}$$

To each such singularity there exists a crepant resolution given by replacing the isolated singular point by a chain of rational curves with intersection pattern dual to the corresponding Dynkin diagram. For example we have

$$Y_{A_n} \rightarrow V(w^2 + x^2 + y^{n+1})$$

where the fibre over $(0, 0, 0)$ consist of n copies of $\mathbb{P}^1$, with the i th curve intersecting the $i + 1$ st at a single point. The threefold $\mathbb{C} \times Y_{A_n}$ is Calabi–Yau and derived equivalent to a quiver with potential. In the A_4 -case the quiver Q_{A_4} is drawn in


FIGURE 1. The quiver Q_{A_4} .

Figure 1. The potential has a similar cubic form to the Hilbert scheme of points considered before

$$W_{A_n} = \left(\sum_{i=0}^{n-1} \alpha_i \right) \cdot \left(\left(\sum_{i=0}^{n-1} \beta_i \right) \cdot \left(\sum_{i=0}^{n-1} \gamma_i \right) - \left(\sum_{i=0}^{n-1} \gamma_i \right) \cdot \left(\sum_{i=0}^{n-1} \beta_i \right) \right)$$

where the above expression is interpreted in the path algebra $\mathbb{C}Q_{A_n}$ of the quiver Q_{A_n} . A similar description exist for the other Lie types above [9]. Continuing with our example, the algebra $A = \mathbb{C}Q_{A_n}/I_{W_{A_n}}$ is three dimensional Calabi–Yau and derived equivalent to the threefold $\mathbb{C} \times Y_{A_n}$. Now one can construct moduli spaces of finite dimensional framed A -modules. By Theorem 3.1 we have (for θ -generic) a description of the moduli space of framed d -dimensional Q_{A_n} -representations in terms of a principal G_d bundle

$$\pi : (H_d \times V_f)^{\theta_{stable}} \rightarrow M_d^{f,\theta}(Q).$$

Then the moduli scheme of stable framed d -dimensional $A = \mathbb{C}Q_{A_n}/I_{W_{A_n}}$ modules is given by the degeneracy locus of the function $\bar{f}_{W_{A_n}} : M_d^{f,\theta}(Q) \rightarrow \mathbb{C}$.

Theorem 1.4. *Let Q, W be the quiver and potential associated to one of the ADE types. Then for θ -generic Behrend’s function is constant on the moduli scheme of d -dimensional θ -stable framed $A = \mathbb{C}Q/I_W$ modules. Moreover we have*

$$\nu_{\{d\bar{f}_W=0\}} = (-1)^{\sum_{i \in I} f_i d_i}$$

where f_i are the components of the framing vector.

Proof. This is identical to the second argument we gave for the Hilbert scheme of points. For generic θ we have a principal G_d bundle $\pi : (H_d \times V_f)^{\theta_{stable}} \rightarrow M_d^{f,\theta}(Q)$ so it suffices to work on the pre-quotient and show the corresponding Milnor fibers are all zero. The reduced variety is now a generalization of the commuting variety but the same argument as in the previous section goes through line by line. The simple sign comes from the symmetry of the quiver. $\square$

It has been shown (see [13]) that for certain choices of stability parameter this quiver perspective produces the familiar moduli spaces of DT/PT objects on $\mathbb{C} \times Y_Q$. In particular in the DT chamber we find the Hilbert scheme of points and curves on $\mathbb{C} \times Y_Q$. As a corollary we have a reducedness statement for the Hilbert scheme of points and curves.

Corollary 1.5. *For all Hilbert polynomials $p(n) = an + b$ the components of $\text{Hilb}^{p(n)}(\mathbb{C} \times Y_Q)$ are generically reduced and equi-dimensional mod 2.*

ACKNOWLEDGEMENTS.

I would like to thank Jim Bryan, Rahul Pandharipande, and Richard Thomas for helpful discussions and comments. I am also grateful to the organizers of the AIM workshop: Motivic Donaldson–Thomas theory and singularity theory for their kind invitation to Budapest. This work was completed at ETH Zürich where I am a postdoc in the research group of Rahul Pandharipande supported by the Swiss research grant SNF 200021.143274.

REFERENCES

- [1] K. Behrend, *Donaldson-Thomas Type Invariants via Microlocal Geometry*, Annals of Math, Pages 1307-1338 from Volume 170 (2009), Issue 3.
- [2] K. Behrend, B. Fantechi *Symmetric obstruction theories and Hilbert schemes of points on threefolds*, Algebra Number Theory 2 (2008) 313-345.
- [3] K. Behrend, J. Bryan, B. Szendrői, *Degree zero motivic Donaldson-Thomas invariants*, arXiv:0909.5088v2.
- [4] J. Denef, F. Loeser, *Geometry on arc spaces of algebraic varieties*, European Congress of Mathematics, Vol. I (Barcelona, 2000), 327348,
- [5] B. Davison, S. Meinhardt, *Motivic DT-invariants for the one loop quiver with potential*, arXiv:1108.5956
- [6] J. Engel, M. Reineke *Smooth models of quiver moduli*, Math. Z. 262 (2009), 4, 817-848.
- [7] W. Fulton, *Intersection Theory*, Springer; 2nd ed. edition.
- [8] L. Göttsche, *Hilbert Schemes of Zero-Dimensional Subschemes of Smooth Varieties*, LNM 1572,. Berlin-Heidelberg-New York 1994.
- [9] A. Gholampour, Y. Jiang, *Counting invariants for the ADE McKay quivers*, 2009, arXiv:0910.5551.
- [10] J. Igusa, *An Introduction to the Theory of Local Zeta Functions*, AMS/IP Studies in Advanced Mathematics.
- [11] A. D. King, *Moduli of representations of finite-dimensional algebras*, Quart. J. Math. Oxford Ser. (2), 45 (1994), no.,180,
- [12] D. Mumford, *Further pathologies in algebraic geometry*, Amer. J. Math. 84 (1962), 642648.
- [13] K. Nagao, H. Nakajima, *Counting invariant of perverse coherent sheaves and its wallcrossing*, International Mathematics Research Notices, vol. 2011 2
- [14] A. Okounkov, N. Reshetikhin, C. Vafa *Quantum Calabi-Yau and Classical Crystals*, arXiv:hep-th/0309208
- [15] R. Pandharipande, A. Pixton, *Gromov-Witten/Pairs correspondence for the quintic 3-fold*, arXiv:1206.5490
- [16] R. Pandharipande, R.P. Thomas, *Curve counting via stable pairs in the derived category*, Inventiones mathematicae, 2009, Vol:178, Pages:407-447
- [17] D. Maulik, N. Nekrasov, A. Okounkov, R. Pandharipande, *Gromov-Witten theory and Donaldson-Thomas theory, I*, Compos. Math. 142 (2006)
- [18] B. Szendrői, *Non-commutative Donaldson-Thomas theory and the conifold*, Geometry & Topology 12 (2008) 11711202. 1171.
- [19] R. P. Thomas, *A holomorphic Casson invariant for Calabi-Yau 3-folds, and bundles on K3 fibrations* J. Differential Geom. Volume 54, Number 2.
- [20] C.M. Walton, *On Degenerations and Deformations of Sklyanin Algebras*, PhD thesis University of Michigan.